SAINT JOHN'S OUTDOOR UNIVERSITY BIRDING LIST—pg. 1

| Common Name | Summer Resident | Migrant (Spring or Fall) | Winter Resident |
|------------------------|--------------------|-----------------------------|--------------------|
| WATERFOWL | | | |
| Snow Goose | | 0 | Sal |
| Canada Goose | С | c (| 0 |
| Trumpeter Swan | | 0 | |
| Tundra Swan | War. | 0 | |
| Wood Duck | C. | C | 115 |
| Gadwall | 12184 | U | й. Х., |
| American Wigeon | 121139 | U | |
| American Black Duck | | | |
| Mallard | С | С | o |
| Blue-winged Teal | C C | c C | 12.0 |
| Northern Shoveler | R | с | 10 |
| Northern Pintail | R | c c | |
| Green-winged Teal | 0 | DC S | |
| Canvasback | 7239 | 0 | 13 |
| Redhead | | U | ų, |
| Ring-necked Duck | 0 | C c c b | 10 |
| Greater Scaup | 160 | R | í. |
| Lesser Scaup | 2 CONSCIENCES | с | |
| Bufflehead | 120 | C A | |
| Common Goldeneye | | 0 | ł |
| Hooded Merganser | U | 6 | 5 |
| Common Merganser | P | 0 | |
| Red-breasted Merganser | 2 | V 0 | |
| Ruddy Duck | | 0 | |
| GROUSE, PTARMIGANS | 5 | 1.1 | 5 |
| Gray Partridge | R | R | R |
| Ring-necked Pheasant | U | j j | U |
| Ruffed Grouse | 0 | 0 | 0 |
| TURKEYS | | - 199 | H X |
| Wild Turkey | С |) c // | C |
| LOON | | 5 12 | VICE. |
| Common Loon | C | С | |
| GREBES | | | |
| Pied-billed Grebe | U | U | |

| V | Common Name | Summer Resident | Migrant (Spring or Fall) | Winter Resident | |
|--------|----------------------------|--------------------|-----------------------------|--------------------|--|
| | Horned Grebe | all and a | 0 | | |
| | Red-necked Grebe | | R | Killer , | |
| | Eared Grebe | | R | 0.918 | |
| | PELICANS | <u> </u> | 次化却 | 1.1 | |
| | American White Pelican | L. XI | 0 | | |
| 37 | CORMORANTS | | | S 41 | |
| | Double-crested Cormorant | | U | S. S. A. | |
| | HERONS | 14.54 | | | |
| | American Bittern | | <u>∫</u>) o < | 16 | |
| Ū | Great Blue Heron | с | - C 🦿 | 107 5 | |
| | Great Egret | 0 | 0 | 12 | |
| | Cattle Egret | 100 | R | | |
| | Green Heron | U | Ú | de- | |
| je | Black-crowned Night-Heron | 0 | 0 | 24 | |
| şĢ | Yellow-crowned Night-Heron | R | R | | |
| T. | VULTURES | | | 825 | |
| | Turkey Vulture | C | C C | Y4 | |
| | HAWKS, EAGLES, & ALLIES | 24 | -27 | 05 | |
| | Osprey | | U | - ~ | |
| | Bald Eagle | U | U | U | |
| | Northern Harrier | Ų | | | |
| | Sharp-shinned Hawk | | U | - | |
| | Cooper's Hawk | U~< | U | | |
| þ | Northern Goshawk | ~/. | R | R | |
| Þ | Red-shouldered Hawk | U/U/ | U | R | |
| | Broad-winged Hawk | /U/ | U | Ker | |
| | Red-tailed Hawk | c | с | Ŭ | |
| | Rough-legged Hawk | ML. |)/ | 0 | |
| | FALCONS | | _r\ # | VIC | |
| | American Kestrel | U | U | KK S | |
| | Merlin | 12 | 0 | | |
| | Peregrine Falcon | 1.22 | 0 | 1.00 | |
| | RAILS, GALLINULES, & COOT | s | | | |
| \Box | Yellow Rail | | R | | |
| \Box | Virginia Rail | | U | | |

| V | Common Name | Summer Resident | Migrant (Spring or Fall) | Winter Resident |
|----|-------------------------|--------------------|--|--------------------|
| | Sora | U | U | |
| | Common Moorhen | R | | |
| | American Coot | U | С | |
| | CRANES | | | |
| | Sandhill Crane | U | U | |
| 62 | PLOVERS | | | |
| | American Golden-Plover | | 0 | |
| | Semipalmated Plover | | U | |
| | Killdeer | d c | С | |
| 52 | SANDPIPERS | Ella | | |
| | Greater Yellowlegs | 1.1 | 0 | |
| | Lesser Yellowlegs | | с | |
| | Solitary Sandpiper | | с | |
| | Spotted Sandpiper | 0 | с | |
| | Upland Sandpiper | 0157 | 0 | |
| | Semipalmated Sandpiper | | t de la constante de la consta | |
| 2Ô | Least Sandpiper | | 2 /c | |
| | Baird's Sandpiper | | Ų | |
| | Pectoral Sandpiper | | C | |
| | Stilt Sandpiper | | 0 | |
| | Buff-breasted Sandpiper | | R | |
| | Short-billed Dowitcher | E.A. | U | |
| | Common Snipe | SA- | с | |
| | American Woodcock | U | U | |
| | Wilson's Phalarope | X | 0 | |
| 1 | | 2E | | |

| | C = Common: Regularly present, easy to find and see in |
|----|--|
| | appropriate habitat at Saint John's. |
| Ц | U = Uncommon: Regularly present in small numbers, |
| ł, | less easy to find and see at Saint |
| 1 | John's. |
| | 0 = Occasional: Irregularly present, a year or two may |
| | elapse between sightings at Saint |
| | John's. |
| | R = Rare: Has occurred at least once at Saint John's |
| | (Outside of normal range). |
| | · · · · · · · · · · · · · · · · · · · |

SAINT JOHN'S OUTDOOR UNIVERSITY BIRDING LIST-pg. 2

| Common Name | Summer Resident | Migrant (Spring or Fall) | Winter Resident | | |
|---------------------------|--------------------|-----------------------------|--------------------|--|--|
| GULLS & TERNS | • | | | | |
| Franklin's Gull | | U | 101 | | |
| Bonaparte's Gull | | 0 | | | |
| Ring-billed Gull | 0 | D U | | | |
| Herring Gull | HALL I | 0 | | | |
| Caspian Tern | | 0 | | | |
| Forster's Tern | | 0 | \mathbf{A}_{i} | | |
| Black Tern | U | Û Û | | | |
| PIGEONS & DOVES | STORY . | 37.72 | | | |
| Rock Pigeon | С | C | С | | |
| Mourning Dove | C C | S c S | -0 | | |
| СИСКООЗ | 22 | | 12 | | |
| Black-billed Cuckoo | 0 | o S | | | |
| Yellow-billed Cuckoo | < U | νiΩu. Š | | | |
| OWLS | 2620 | Xee. | 2.1 | | |
| Eastern Screech-Owl | 0 | 0 | 0 | | |
| Great Horned Owl | 0 | ~ os | 0 | | |
| Snowy Owl | 466 | -NSK | R | | |
| Barred Owl | C U C | NU. | U | | |
| Great Gray Owl | 130 | | R | | |
| Long-eared Owl | R | R | R | | |
| Short-eared Owl | | 0 | 5 | | |
| Boreal Owl | P / | 0 | R | | |
| Northern Saw-whet Owl | 0. | 0 | R | | |
| NIGHTJARS | -2 | | e.K | | |
| Common Nighthawk | U | с | 5 | | |
| Whip-poor-will | | 0 | 62 | | |
| SWIFTS | | - Ks | 6.728 | | |
| Chimney Swift | С | С | 1 3 | | |
| HUMMINGBIRDS | | 19 | 11 2 | | |
| Ruby-throated Hummingbird | С | . c // | | | |
| KINGFISHERS | | S. H. | | | |
| Belted Kingfisher | U* | U | | | |
| WOODPECKERS | | | | | |
| Red-headed Woodpecker | 0 | U | R | | |

| 1 | Common Name | Summer Resident | Migrant (Spring or Fall) | Winter Resident |
|--------|---------------------------|---|-----------------------------|--------------------|
| | Red-bellied Woodpecker | Sho . | C, | U |
| | Yellow-bellied Sapsucker | U | u V | Killer I |
| | Downy Woodpecker | С | С | с |
| | Hairy Woodpecker | с | C L | с |
| | Black-backed Woodpecker | | 400 | R |
| | Northern Flicker | С | C / | R |
| | Pileated Woodpecker | c | - c | с |
| R | FLYCATCHERS | | | |
| | Olive-side Flycatcher | Į, | ○] 0 ≤ | 100 |
| | Eastern Wood-Pewee | c | - c 🦾 | RUT S |
| | Yellow-bellied Flycatcher | 1897 | 0 | 12 |
| | Alder Flycatcher | | 0 | |
| | Willow Flycatcher | 0 | 0 | de- |
| | Least Flycatcher | c | C | 34 |
| ģ | Eastern Phoebe | с | ¢ | |
| | Great Crested Flycatcher | с | с | Å. |
| | Western Kingbird | | R | -74 |
| \Box | Eastern Kingbird | C | C | |
| | SHRIKES | | | e 7-4 |
| | Northern Shrike | | 0 | 0 |
| | VIREOS | ~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~ | 4 | |
| | Yellow-throated Vireo | | С | |
| | Blue-headed Vireo | | U | |
| Ó | Warbling Vireo | U/ | 0 | |
| Ó | Philadelphia Vireo | NG / / | 0 | |
| | Red-eyed Vireo | C | C | nt - |
| U | JAYS, CROWS, & ALLIES | W/A | N. | FF |
| | Gray Jay | | 1 | R |
| | Blue Jay | с | С | / U |
| | American Crow | С | C C | U T |
| | Clark's Nutcracker | | 10X "1 | \mathcal{S} |
| | LARKS | 1.428 | 1.56.0 | 1.20 |
| | Horned Lark | 0 | Ŭ | 0 |
| | SWALLOWS | | | |
| | Purple Martin | С | С | |

| \checkmark | Common Name | Summer Resident | Migrant (Spring or Fall) | Winter Resident | |
|--------------|-------------------------------|--------------------|-----------------------------|--------------------|--|
| | Tree Swallow | С | С | | |
| | Northern Rough-winged Swallow | U | U | | |
| | Bank Swallow | U | U | | |
| | Cliff Swallow | С | С | | |
| | Barn Swallow | С | С | | |
| Ð, | CHICKADEES | | | | |
| | Black-capped Chickadee | С | С | С | |
| | Boreal Chickadee | | | R | |
| | NUTHATCHES | | | | |
| | Red-breasted Nuthatch | 0 | U | U | |
| | White-breasted Nuthatch | С | С | С | |
| 2 | CREEPERS | | | | |
| | Brown Creeper | 0 | U | 0 | |
| 4 | WRENS | 42 G | | | |
| | House Wren | с | Ċ | | |
| 口 | Winter Wren | | P | | |
| ø | Sedge Wren | U | 2 | | |
| | Marsh Wren | - U | U | | |
| Z | KINGLETS | | | | |
| | Golden-crowned Kinglet | | U | 0 | |
| | Ruby-crowned Kinglet | | U | 0 | |
| | Blue-gray Gnatcatcher | U . | U | | |
| | STARLINGS | SH - | - | | |
| \Box | European Starling | С | С | U | |

C = Common: Regularly present, easy to find and see in appropriate habitat at Saint John's. U = Uncommon: Regularly present in small numbers, less easy to find and see at Saint John's. O = Occasional: Irregularly present, a year or two may elapse between sightings at Saint John's. R = Rare: Has occurred at least once at Saint John's (Outside of normal range).

SAINT JOHN'S OUTDOOR UNIVERSITY BIRDING LIST-pg. 3

Last Updated: March 2020

| Common Name | Summer Resident | Migrant (Spring or Fall) | Winter Resident | Common Name | Summer Resident | Migrant (Spring or Fall) | Winter Resident | \checkmark | Common Name | Summer Resident | Migrant (Spring or Fall) | Winter Residen |
|------------------------------|--------------------|-----------------------------|--|---------------------------|---|-----------------------------|--|--------------------------------|---|--------------------|-----------------------------|-------------------|
| THRUSHES | | | Black-and-White Warbler | Black-and-White Warbler C | | | | CARDINALS, GROSBEAKS, & ALLIES | | | | |
| Eastern Bluebird | С | С | R - 30 | American Redstart | С | Ċ | Care . | | Northern Cardinal | С | С | С |
| Veery | U | υ, (, | A 182 | Prothonotary Warbler | | R | 1. 11/2 | | Rose-breasted Grosbeak | С | С | |
| Gray-cheeked Thrush | | 0 | | Ovenbird | C | с | 13. 21 | | Indigo Bunting | U | U | |
| Swainson's Thrush | in the second | С | | Northern Waterthrush | L > M | J U | 4474 | 97 | Dickcissel | | 0 | |
| Hermit Thrush 💦 👌 | はわろか | С | | Louisiana Waterthrush | Kod A | R | 5 5 1 | | BLACKBIRDS, ORIOLES, & AL | LIES | | |
| Wood Thrush | 0 | о — | | Connecticut Warbler | | 0 | 550 | | Bobolink | | 0 | |
| American Robin | С | С | U | Mourning Warbler | 0 | Ċ | L KK | | Red-winged Blackbird | С | С | 0 |
| MIMIC THRUSHES | 217 655 | ê. 93 | | Common Yellowthroat | C | C C | 57.0 | | Eastern Meadowlark | 0 | 0 | |
| Gray Catbird | с | с | DA. | Wilson's Warbler | | C di | 5007 (= | | Western Meadowlark | U U | U | |
| Northern Mockingbird | 112 | 0 | $(\mathcal{L}_{\mathcal{M}})$ | Canada Warbler | 1897 | U | 11 - 11 | | Yellow-headed Blackbird | U | U | |
| Brown Thrasher | U | U | | Prairie Warbler | 1947 - 1949 - 1949 - 1949 - 1949 - 1949 - 1949 - 1949 - 1949 - 1949 - 1949 - 1949 - 1949 - 1949 - 1949 - 1949 - | R | K [24 | | Rusty Blackbird | | U | R |
| PIPITS | | 2 Y. | | Blackpoll Warbler | 4.62 | U. | | 90 | Brewer's Blackbird | | 🖉 о | |
| American Pipit | | 0 | | TANAGERS | | 1000 | and a | Ó | Common Grackle | с | с | |
| WAXWINGS | 1755 | | | Scarlet Tanager | U (| (7 U) | | | Brown-headed Cowbird | C - | с | |
| Bohemian Waxwing | | 0 | 0 | SPARROWS | 35 | | 201 | | Orchard Oriole | 0 | 0 | |
| Cedar Waxwing | C > | c | U Lie | Eastern Towhee | | 0 | TLA | R | Baltimore Oriole | с | -/c | |
| WARBLERS | | -SAK | 6 | American Tree Sparrow | | c | | Ē | FINCHES | | | |
| Blue-winged Warbler | R | R | | Chipping Sparrow | С | C | | | Pine Grosbeak | | 0 | 0 |
| Golden-winged Warbler | R | 0 | | Clay-colored Sparrow | 0 | U | | | Purple Finch | 0 | υ | 0 |
| Tennessee Warbler | | с | λ | Field Sparrow | 0 | 4,0 | | | | с | с | U |
| Orange-crowned Warbler | Ű | U C | S | Vesper Sparrow | 0 | U | | | Red Crossbill | 0 | 0 | 0 |
| Nashville Warbler | 0 | С | | Lark Sparrow | | R | | | White-winged Crossbill | 546 | 0 | 0 |
| Northern Parula | 0.5 | U | | Savannah Sparrow | | U | | | Common Redpoll | | 0 | U |
| Yellow Warbler | - Yc | с | al | Fox Sparrow | N/I | U | | \Box | Pine Siskin | R | 0 | 0 |
| Chestnut-sided Warbler | U | с | - 5 | Song Sparrow | c | C, | | | American Goldfinch | c | С | U |
| Magnolia Warbler | | U | all a | Lincoln's Sparrow | 11/- | U | FE | | Evening Grosbeak | | 0 | 0 |
| Cape May Warbler | 7 | Ů. | | Swamp Sparrow | U | С | V_{1} | \Box | WEAVERS | | | |
| Yellow-rumped Warbler | | C | N. | White-throated Sparrow | 16 | c | R | la. | House Sparrow | С | С | C |
| Black-throated Green Warbler | | <u>ں</u> | 4 2 | Harris's Sparrow | 11 · · | a Will | 16 3 | - 6 | C = Common: Regularly pres | ent, easy to | o find and se | ee in |
| Blackburian Warbler | | ο U | \owned to be a construction of the constructio | White-crowned Sparrow | 10 | U | | \mathbf{F}_{1} | appropriate ha | bitat at Sai | int John's. | |
| Pine Warbler | | JU // | | Dark-eyed Junco | 1000 | с | С | | U = Uncommon: Regularly p easy to find | | | |
| Palm Warbler | · · · · · · · · | U | | Lapland Longspur | | 0 | | | 0 = Occasional: Irregularly p | resent, a ye | ear or two n | nay |
| Bay-breasted Warbler | | U | | Snow Bunting | | | | | elapse betwe | en sighting | s at Saint J | ohn's. |
| Cerulean Warbler | U | U | | | | | <u>. </u> | | R = Rare: Has occurred at lea (Outside of normal | | Saint John | s |